



## **Remasan Product Details**

### **Description**

REMASAN® Globuli



### **Active & Inactive Ingredients**

Active Ingredients:

1 Globulus containing 200mg contains: Arnica montana C12, C30 and C200

Inactive Ingredients: 200mg Saccharose

### **Dosage & Form**

Globules

Appearance: white saccharose pellets with a diameter of approximately 6mm, and a weight of approximately 0,2g in a blister package.

### **Clinical Specifications**

#### **Area of application:**

Pain reliever & Trauma remedy.

This includes acute injuries of all kinds, contusions, cuts, bruises, (bleeding) wounds and sport injuries, nose bleeds.

Chronic complaints due to the above-mentioned occurrences such as painful conditions or mood disorders after injury.

Pre- and post- surgery;

The use of these homeopathic remedies in the afore mentioned areas are based solely on experience with homeopathic remedies.

In severe illnesses, homeopathic remedies should not replace clinically proven treatment.

#### **Dosage, form and duration of use;**

*Acute injuries:*

REMASAN should be taken as soon as possible after injury.



When taken, the pellets should be allowed to dissolve under the tongue. No food or drink should be consumed immediately before or after the remedy has been taken.

In case the effect wears off, the procedure may be repeated.

*Chronic ailments:*

Globules can be taken 2x daily (melt under tongue).

*Pre and post operations:*

Let a globule melt under the tongue, 1x daily, beginning 1 day before, and 3 days after surgery.

*Dosage for children:*

Dosage and duration of use is the same as with adults.

Treatment of infants younger than one year old, must be discussed with a doctor.

## **Warnings**

In case of an accident, first aid measures for emergencies must be administered before taking Remasan. Please ask a doctor or pharmacist in case the pain continues for a prolonged time.

Do not take Remasan Globules in case you suffer from sugar intolerance.

Patients with rare genetic fructose and galactose intolerance, a glucose malabsorption or a sucroseisomaltase insufficiency should not use this medicine.

Reference for diabetics: one globule contains 0,2g of saccharose.

Contraindication with other remedies and other reactions There have been no studies on this subject.

Remasan is safe if pregnant or breast-feeding.

A negative influence on driving or an incapacity to operate machines has not been reported yet.

Side Effects: No side effects up to date.

Overdose: No reports of an OD up to date.

## **Pharmaceutical characteristics**

Inactive ingredients: Saccharose

Incompatibility: Not-applicable

Expiration Date: 5 years after production

Remedia Homöopathie  
Mag. pharm. Robert Müntz GesmbH  
Hauptstraße 4  
A-7000 Eisenstadt



Safety measures for storage:

Do not store over 25°C (77°F); do not store near electro-magnetic fields (eg. Mobile phones, TV, computers, microwave, etc.).

Type and content of the container: Blisterpack of 10 or 30 globules size 8

Particular safety measures for disposal or other handling: No particular measures necessary.

### **Licensed Owner**

Remedia Homöopathie  
Mag.pharm. Robert Müntz GesmbH  
Hauptstraße 4, 7000 Eisenstadt  
Tel.: 02682 62220 66  
Fax: 02682 62220 62  
e-Mail: [info@remedia.at](mailto:info@remedia.at)

### **Registration Number**

3-00334

**Orders can be placed online: [www.remedia.at](http://www.remedia.at)**